arXiv:nlin/0609019v2 [nlin.SI] 13 Sep 2006

A note on elliptic coordinates on the Lie algebra $e(3)$.

A.V. Tsiganov
St.Petersburg State University, St.Petersburg, Russia
e–mail: tsiganov@mph.phys.spbu.ru

Abstract

We introduce elliptic coordinates on the dual space to the Lie algebra $e(3)$ and discuss the separability of the Clebsch system in these variables. The proposed Darboux coordinates on $e^*(3)$ coincide with the usual elliptic coordinates on the cotangent bundle of the two-dimensional sphere at the zero value of the corresponding Casimir function.

The Lie algebra $e(3) = so(3) \oplus \mathbb{R}^3$ of the Lie group of euclidean motions of $\mathbb{R}^3$ is a semidirect sum of an algebra $so(3)$ and an abelian ideal $\mathbb{R}^3$. For convenience we shall use the invariant inner product to identify the dual of the Lie algebra, namely $e^*(3)$, with the Lie algebra $e(3)$.

On the dual space $e^*(3)$ with coordinates $J = (J_1, J_2, J_3) \in so(3) \simeq \mathbb{R}^3$ and $x = (x_1, x_2, x_3) \in \mathbb{R}^3$ the Lie-Poisson bracket is defined by

$$\{J_i, J_j\} = \varepsilon_{ijk} J_k, \quad \{J_i, x_j\} = \varepsilon_{ijk} x_k, \quad \{x_i, x_j\} = 0\,, \tag{1}$$

where ε_{ijk} is the sign of the permutation (ijk) of (123). The Lie-Poisson bracket (1) is degenerated and has two Casimir functions

$$A = |x|^2 \equiv \sum_{k=1}^{3} x_k^2, \qquad B = \langle x, J\rangle \equiv \sum_{k=1}^{3} x_k J_k. \tag{2}$$

Here $\langle x, J\rangle$ stands for the standard euclidean scalar product in $\mathbb{R}^3$. The generic symplectic leaf

$$\mathcal{O}_{ab} = \{x, J\,:\; A = a^2, \;\; B = b\}\,, \tag{3}$$

is a four-dimensional symplectic manifolds, which is topologically equivalent to the cotangent bundle $T^*\mathbb{S}^2$ of the two-dimensional sphere $\mathbb{S}^2 = \{x \in \mathbb{R}^3, |x| = a\}$ [1].

If $b = 0$, then there exists a symplectic transformation

$$\rho : (p, x) \to J = p \wedge x, \qquad J_i = \sum_{j,k=1}^{n=3} \varepsilon_{ijk}\, p_j\, x_k\,, \tag{4}$$

which identify $T^*\mathbb{S}^2 \subset T^*\mathbb{R}^3$ and $\mathcal{O}_{a0}$. Here $p \wedge x$ means the standard euclidean cross product in $\mathbb{R}^3$ and vector p is canonically conjugated to x momenta in $T^*\mathbb{R}^3$, $\{p_i, x_j\} = \delta_{ij}$, such that $\langle p, x\rangle = 0$.

If $b \neq 0$, the symplectic structure of manifold $\mathcal{O}_{ab}$ differs from the standard symplectic structure of $T^*\mathbb{S}^2$ by a magnetic term proportional to b [1].

The aim of this note is to describe this magnetic term with the help of elliptic coordinates on the sphere $\mathbb{S}^2$ lifted to the Darboux variables on the manifold $e^*(3)$ by $b \neq 0$.

The elliptic coordinates u_1, u_2 on $\mathbb{S}^2$ with parameters $\alpha_1 < \alpha_2 < \alpha_3$ is defined as roots of the equation

$$e(\lambda) = \sum_{j=1}^{3} \frac{x_j^2}{\lambda - \alpha_j} = \frac{a^2(\lambda - u_1)(\lambda - u_2)}{\varphi(\lambda)} = 0, \tag{5}$$

where $\varphi(\lambda) = \prod_{j=1}^{3}(\lambda - \alpha_j)$ and $|x| = a$, see [2]. Like the elliptic coordinates in $\mathbb{R}^3$, the elliptic coordinates on $\mathbb{S}^2$ are also orthogonal and only locally defined. They take values in the intervals

$$\alpha_1 < u_1 < \alpha_2 < u_2 < \alpha_3. \tag{6}$$

By using the Lie-Poisson bracket (1) we can prove that elliptic coordinates $u_{1,2}$ and variables

$$\pi_{1,2}^0 = h(u_{1,2}), \qquad h(\lambda) = \frac{1}{2a^2} \sum_{j=1}^{3} \frac{x_j\,(x \wedge J)_j}{\lambda - \alpha_j} \tag{7}$$

satisfy to the following relations

$$\{u_1, u_2\} = 0, \qquad \{\pi_i^0, u_j\} = \delta_{ij}, \qquad \{\pi_1^0, \pi_2^0\} = \frac{ib}{4a}\,\frac{u_2 - u_1}{\sqrt{\varphi(u_1)\varphi(u_2)}}. \tag{8}$$

These relations may be easy obtained by means of the Poisson brackets between the generating functions $e(\lambda)$ and $h(\mu)$

$$\begin{aligned}
\{e(\lambda), e(\mu)\} &= 0, \qquad \{e(\lambda), h(\mu)\} = -a^{-2} e(\lambda)\, e(\mu) - \frac{e(\lambda) - e(\mu)}{\lambda - \mu},\\
\{h(\lambda), h(\mu)\} &= \frac{b}{4a^2}\,\frac{x_1 x_2 x_3 (\lambda - \mu)(\alpha_1 - \alpha_2)(\alpha_2 - \alpha_3)(\alpha_3 - \alpha_1)}{\varphi(\lambda)\,\varphi(\mu)}.
\end{aligned}$$

If $b = 0$ relations (8) yield well known fact that variables $u_{1,2}$ and $\pi_{1,2}^0$ are the Darboux coordinates on the manifold $T^*\mathbb{S} \simeq \mathcal{O}_{a0}$.

The coordinates u_i and the parameters α_j can be subjected to a simultaneous linear transformation $u_i \to \beta u_i + \gamma$ and $\alpha_j \to \beta\alpha_j + \gamma$, so it is always possible to choose

$$\alpha_1 = 0, \qquad \alpha_2 = 1, \qquad \alpha_3 = \mathtt{k}^2 > 1.$$

Using relations (8) and this choice of parameters α_j we can prove the following

Proposition 1 *If $b \neq 0$, elliptic coordinates $u_{1,2}$ (5) and the corresponding momenta*

$$\pi_{1,2} = \pi^0_{1,2} - bf_{1,2}, \qquad f_{1,2} = \frac{u_{1,2}}{2a\sqrt{\varphi(u_{1,2})}}\, F\left(\frac{\sqrt{\mathtt{k}^2 - u_{2,1}}}{\mathtt{k}}, \frac{\mathtt{k}}{\mathtt{k}^2 - 1}\right), \tag{9}$$

form a complete set of the Darboux variables on the manifold $e^*(3)$

$$\{u_1, u_2\} = \{\pi_1, \pi_2\} = 0, \qquad \{\pi_i, u_j\} = \delta_{ij}\,,$$

which are real variables in their domain of definition (6). Here $F(z,k)$ is incomplete elliptic integral of the first kind, which is identical to the inverse function of the elliptic Jacobi function $sn(z,k)$ [3].

Proof: The functions $f_{1,2}$ depend on the coordinates $u_{1,2}$ and, therefore, we have to verify one relation only

$$\{\pi_1, \pi_2\} = \{\pi^0_1, \pi^0_2\} - b\{\pi^0_1, f_2\} + b\{\pi^0_2, f_1\} = \{\pi^0_1, \pi^0_2\} - b\left(\frac{\partial f_2}{\partial u_1} - \frac{\partial f_1}{\partial u_2}\right) = 0\,.$$

This relation follows from (8) and properties of the incomplete elliptic integral of the first kind $F(z,k)$.

So, equations (5), (7) and (9) define elliptic variables u and π on $e^*(3)$ as functions on initial variables x and J. The inverse transformation $(u,\pi) \to (x,J)$ reads as

$$x_j = a\sqrt{\frac{(\alpha_j - u_1)(\alpha_j - u_2)}{(\alpha_j - \alpha_m)(\alpha_j - \alpha_n)}}\,, \qquad J_j = a^{-2}\Big(bx_j + (z \wedge x)_j\Big), \tag{10}$$

where $m \neq j \neq n$ and entries of the vector z are given by

$$z_j = \frac{2\,x_j}{u_1 - u_2}\left(\frac{\varphi(u_1)(\pi_1 + bf_1)}{\alpha_j - u_1} - \frac{\varphi(u_2)(\pi_2 + bf_2)}{\alpha_j - u_2}\right)\,.$$

Like the elliptic coordinates in $\mathbb{R}^n$ and on $\mathbb{S}^n$ [2], the elliptic variables on $e^*(3)$ may be successfully exploited in the theory of integrable systems. For instance, substituting expressions (10) into the quadratic hamiltonian

$$H = \frac{1}{2}(J_1^2 + J_2^2 + J_3^2) + \frac{1}{2}(\alpha_1 x_1^2 + \alpha_2 x_2^2 + \alpha_3 x_3^2),$$

associated with the integrable Clebsch system on $e^*(3)$, one gets this hamiltonian in terms of the elliptic variables

$$H = \frac{2}{u_1 - u_2}\Big(\varphi(u_1)(\pi_1 + bf_1)^2 - \varphi(u_2)(\pi_2 + bf_2)^2\Big) + \frac{b^2}{2a^2} - \frac{a^2(u_1 + u_2 - \sum_{j=1}^3 \alpha_j)}{2}\,.$$

Here $f_{1,2}$ are functions on u_1 and u_2 and, therefore, the hamiltonian H belongs to the Stäckel family of integrals of motion at $b = 0$ only, i.e. for the Neumann system on the sphere [2]. In spite of the fact that $f_{1,2}$ are elliptic functions this hamiltonian H may be rewritten as rational quasi-Stäckel hamiltonian introduced in [4].

Thus we have to fully appreciate that elliptic variables $u_{1,2}$ and $\pi_{1,2}$ cannot be the separation variables for the Hamilton-Jacobi equation associated with the Clebsch system at $b \neq 0$. The similar result has been obtained in [5, 6] by using velocities

$$\dot{u}_{1,2} = \frac{\mp 4\varphi(u_{1,2})(\pi_{1,2} + bf_{1,2})}{u_1 - u_2}$$

instead of the momenta $\pi_{1,2}$. It will be interesting to compare this result with results of the paper [7] about separability of the Clebsch system by using the "Kowalevski variables", which in fact coincide with the elliptic coordinates (5).

On the other hand, we can substitute Darboux variables $u_{1,2}$ and $\pi_{1,2}$ into the usual Stäckel integrals of motion and get a whole family of integrable systems on the manifold $e^*(3)$, which are separable in these variables. The main problem in this widely known Jacobi method is how to single out integrable systems interesting in physics from this huge family.

The research was partially supported by the RFBR grant 06-01-00140. The author want to thank V.G. Marikhin for valuable discussions.

References

[1] S.P. Novikov, *Hamiltonian formalism and multi-valued analog of Morse theory*, Uspekhi Mat. Nauk, **37**, 3-49, 1982.

[2] E. G. Kalnins, *Separation of Variables for Riemannian Spaces of Constant Curvature*, Longman Scientific & Technical, Essex, 1986.

[3] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, Dover Publications Inc., New York, 1046 p., 1965.

[4] V.G. Marikhin, V.V. Sokolov, *On the quasi-Stäckel hamiltonians*, Uspekhi Mat. Nauk, **60**(5), 175-176, 2006.

[5] E.I. Kharlamova, *O dvigenii tverdogo tela vokrug nepodvignoj tochki v tsentral'nom n'yutonovskom pole*, Izvestiya Sib. Otd. AN SSSR, **6**, 7-17, 1959.

[6] I.V. Komarov, A.V. Tsiganov, *On a trajectory isomorphism of the Kowalevski gyrostat and the Clebsch problem*, J. Phys.A, **38**, 2917-2927, 2005.

[7] V.G. Marikhin, V.V. Sokolov, *Separation of variables on a non-hyperelliptic curve*, Reg. Chaot. Dyn, **10**(1), 59-70,2005.